

Subscribe to the SAJAA and earn 15 CPD points per year

CPD reference number A008/016/03/2002

Continuing Professional Development Point Questionnaire

Accredited by the Wits CPD Office



- Sepsis is the systemic inflammatory response to infection.
 (a) true
 (b) false
- 2. Sepsis is caused by the activation of a single gene.
 - (a) true (b) false
- Hypotension in septic shock is due increased vascular tone.
 (a) true (b) false
- 4. In Sepsis surgery is an emergency not urgency.
 - (a) true (b) false
- Community acquired sepsis should be treated with new generation antibiotics.
 - (a) true (b) false
- The coagulation cascade is activated by cytokines.
 - (a) true (b) false
- 7. Pressure limit ventilation minimizes respiratory morbidity.
 - (a) true (b) false
- Adrenaline is being used more often as the first line inotrope in septic shock.
 - (a) true (b) false
- With neuropathic pain, the afferent tracts are in normal physiologic status.
 - (a) true (b) false
- 10. Allodynia has the effects of "pain on a touch" stimulus.
 - (a) true (b) false
- Neuropathic pain usually derives from Ad-fibre mal-conduction.
 (a) true (b) false
- "Wind-up" happens in the dorsal horn with WDR cells as major component of activity.
 - (a) true (b) false
- Glutamate is the most important neuro-transmitter in the dorsal horn with respect to the NMDA-receptor.
 - (a) true (b) false
- When compared to CABG, angioplasty has a lower initial cost and results in longer lasting relief of symptoms.
 - (a) true (b) false
- 15. During OPCAB grafting, occluding a severely stenosed vessel (>90%) may lead to less severe myocardial ischaemia than a vessel with only 60-70% stenosis.
 - (a) true (b) false

- Continuous cardiac output monitoring will reflect a change in cardiac output immediately.
 - (a) true (b) false
- Atrial contraction may contribute to up to 50% of LV stroke volume during OPCAB surgery.
 - (a) true (b) false
- Ischaemic pre-conditioning is more powerful than pharmacological cardiac protection.
 - (a) true (b) false
- Children may become agitated after the administration of Midazolam.
 - (a) true (b) false
- 20. Chloral hydrate may cause airway obstruction in children.
 - (a) true (b) false
- 21. EMLA cream has very few side-effects.
 - (a) true (b) false
- 22. Nitrous oxide may cause nausea and vomiting.
 - (a) true (b) false
- 23. Ketamine may be given rectally.
 - (a) true (b) false

Instructions

- 1. Read the journal to find all the answers
- 2. Answer the questions and mark the answer form by putting an "X" in the appropriate square.
- 3. Make a photocopy for your records in case the form is lost in the mail.
- 4. Place the answer form in a envelope, seal it and send to:-

In House Publications SAJAA CPD Points PO Box 412748 Craighall

- 2024
- 5. Do not register it and do not fax it, only original answer sheets are accepted.
- 6. To get your CPD points subscribe to the SAJAA. See subscription form.